The Matrix Code

How the Heisenberg Uncertainty Principle Proves Reality is a Simulation

Rohan Verma, MSc Physics
Indian Institute of Technology Kanpur, India

Preface

This book is the consequence of going on the path of seeking truth or gaining a deeper and deeper understanding of the cosmos and life. It is written for everyone, regardless of their background.

Through this book, I aim to provide insights into how we exist within a simulation or as part of a simulated reality. Given the infinite nature of the cosmos and life, no matter how much we discover, there will always be something beyond. There is no theory, scientific framework, or documented text that can be considered absolutely true or false. Every piece of knowledge ever created is an attempt to understand life and the cosmos to some extent.

If you wish to connect with me for any reason, you can reach me at **rohanv.i341@gmail.com**.

Contents

1 What is the Matrix? 1

2 How Images are Just Matrices 3

3 What is the Heisenberg Uncertainty Principle? 4
3.1 Uncertainty in Time and Energy 4
3.2 Robertson-Schrödinger uncertainty relation 5

4 Phase Space is a Matrix 6
4.1 Phase Space Coordinates 6
4.1.1 Classical Mechanics 6
4.1.2 Dynamical Systems 6
4.1.3 Statistical Mechanics 6
4.1.4 Quantum Mechanics 7
4.1.5 Relativity Cosmology 7
4.1.6 Quantum Field Theory String Theory 7
4.2 Phase space of various systems 8
4.3 Discretization Limit of Phase Space: Emergence of Matrix Structure 13
4.4 Absolute Vacuum or Nothingness Is a Matrix 15
4.5 List of other clues for simulation or matrix 16
4.5.1 Position-Momentum (x, p) 16
4.5.2 Energy-Time (t, E) 16
4.5.3 Angular Position-Angular Momentum (θ, L_z) 16
4.5.4 Number-Phase (N, ϕ) 16

5 Video Game vs. Our Universe 17
5.1 4K Video Game: A Simulation Created by Humans 17
5.1.1 Resolution or Size of the Matrix 17
5.1.2 How the Image Matrix Changes Over Time 17
5.1.3 Powered By 17
5.2 Our Universe 17
5.2.1 Resolution or Size of the Matrix 17
5.2.2 How the Cosmic Matrix Changes Over Time 18
5.2.3 Powered By 20

Chapter 1

What is the Matrix?

A matrix can represent anything—whether it is a physical system, a person, a concept, or even the universe itself.

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \tag{1.1}$$

The matrix **A**, as defined in equation (1.1), is a two-dimensional matrix, where each element (or entity) is one-dimensional.
In the most general form:

- A matrix can exist in n-dimensions, where n can be any positive integer, including infinity.
- An element of a matrix can also have any dimension, including infinity.

Examples of Matrices

1. Matrix Dimension = 1, Element Dimension = 1
   A segment of a DNA sequence can be represented as a one-dimensional matrix, where each base is encoded as a number:

   $$A = 1, \quad T = 2, \quad G = 3, \quad C = 4$$

   Thus, a DNA sequence D can be written as:

   $$D = \begin{bmatrix} 1 & 3 & 4 & 2 & 1 & 3 & 4 \end{bmatrix}$$

2. Matrix Dimension = 2, Element Dimension = 1
   The temperature at different (x, y) locations on a metal plate can be represented using a 2D matrix, where each element is a scalar temperature value.

   $$T = \begin{bmatrix} 300 & 320 & 310 \\ 280 & 290 & 295 \\ 260 & 270 & 275 \end{bmatrix} \text{ K}$$

3. Matrix Dimension = 3, Element Dimension = 3

   In fluid dynamics, the velocity field in three-dimensional space is represented as a 3D matrix (tensor). Each element of this matrix is itself a vector (3D entity) containing velocity components in the x, y, z directions.

   The velocity tensor V can be written as:

$$V = \begin{bmatrix} (v_{111}, v_{112}, v_{113}) & (v_{121}, v_{122}, v_{123}) & \dots \\ (v_{211}, v_{212}, v_{213}) & (v_{221}, v_{222}, v_{223}) & \dots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

   Each element v_{ijk} represents a velocity component in a particular direction at a given spatial coordinate.

Chapter 2

How Images are Just Matrices

An image is a two-dimensional matrix, where each element of the matrix is a three-dimensional vector representing the intensity or amount of Red (R), Green (G), and Blue (B). The combined effect of these color intensities determines the final color of each matrix element.
Each element of the matrix is called a pixel. The resolution of an image defines the total number of pixels present in it.

For example, consider a Full HD image. It has a resolution of 1920 $\times$ 1080 pixels. The matrix is of order 1080 $\times$ 1920.

A 1920 $\times$ 1080 image can be represented as:

$$I = \begin{bmatrix} (R_{1,1}, G_{1,1}, B_{1,1}) & (R_{1,2}, G_{1,2}, B_{1,2}) & \dots & (R_{1,1920}, G_{1,1920}, B_{1,1920}) \\ (R_{2,1}, G_{2,1}, B_{2,1}) & (R_{2,2}, G_{2,2}, B_{2,2}) & \dots & (R_{2,1920}, G_{2,1920}, B_{2,1920}) \\ \vdots & \vdots & \ddots & \vdots \\ (R_{1080,1}, G_{1080,1}, B_{1080,1}) & (R_{1080,2}, G_{1080,2}, B_{1080,2}) & \dots & (R_{1080,1920}, G_{1080,1920}, B_{1080,1920}) \end{bmatrix}$$

Each entry (R_{ij}, G_{ij}, B_{ij}) represents the intensity values of red, green, and blue for the pixel at row i and column j.

Static Images vs. Videos

If the image matrix remains constant over time, it represents a static image. However, if the image matrix changes over time, it forms a video.

Chapter 3

What is the Heisenberg Uncertainty Principle?

The Heisenberg Uncertainty Principle states that it is impossible to simultaneously measure the exact position and momentum of a particle. Mathematically, it is expressed as:

$$\Delta x \cdot \Delta p \geq \frac{\hbar}{2} \approx 5.27286 \times 10^{-35} \text{ J} \cdot \text{s} \tag{3.1}$$

where:

- Δx is the uncertainty in position,
- Δp is the uncertainty in momentum,
- $\hbar$ is the reduced Planck's constant ($\hbar = \frac{h}{2\pi}$).

Example: Electron Diffraction Through a Narrow Slit

A beam of electrons passing through a narrow slit forms a diffraction pattern. The narrower the slit (Δx), the greater the spread in momentum (Δp_x), broadening the diffraction pattern.

- When slit width is reduced, the diffraction pattern widens.
- This matches the uncertainty relation:

$$\Delta x \cdot \Delta p_x \geq \frac{\hbar}{2}$$

3.1 Uncertainty in Time and Energy

The energy-time uncertainty principle states that the uncertainty in energy (ΔE) and the uncertainty in time (Δt) are fundamentally related by:

$$\Delta E \cdot \Delta t \geq \frac{\hbar}{2}$$

This implies that the shorter the time interval over which an event occurs, the greater the uncertainty in its energy.

Example: Lifetime of Excited States in Atoms
An atom in an excited state with energy E decays to a lower level by emitting a photon of energy E_{photon}. The excited state's lifetime (Δt) determines the energy uncertainty (ΔE) of the photon.

- If the excited state has a short lifetime (Δt is small), the emitted photon's energy has greater uncertainty (ΔE is large).
- Conversely, if the excited state has a longer lifetime, the emitted photon's energy has less uncertainty (ΔE is small).

3.2 Robertson-Schrödinger uncertainty relation

$$\Delta A \cdot \Delta B \geq \frac{1}{2} \left| \langle [A, B] \rangle \right|$$

where $\langle [A, B] \rangle$ represents the expectation value of the commutator of A and B.

Chapter 4

Phase Space is a Matrix

Phase space is a mathematical space in which all possible states of a system are represented, with each state corresponding to a unique point in this space.

4.1 Phase Space Coordinates

4.1.1 Classical Mechanics

- Phase space coordinates: (q, p), where q is position and p is momentum.
- A system with N degrees of freedom has a $2N$-dimensional phase space.
- Example: A simple pendulum is described by (θ, p_θ).

4.1.2 Dynamical Systems

- Generalized state variables $(x_1, x_2, \ldots, x_n)$ describe system evolution.
- Example: The Lotka-Volterra system uses $(N_{\text{prey}}, N_{\text{predator}})$.

4.1.3 Statistical Mechanics

- Microcanonical ensemble: Fixed energy surface in (q, p)-space.
- Canonical ensemble: Probability distributions over (q, p) considering temperature.
- Grand canonical ensemble: Includes particle number N, so phase space is (q, p, N).
- Example: An ideal gas system with (q_i, p_i) for each particle.

4.1.4 Quantum Mechanics

- Uses wavefunctions in Hilbert space instead of classical (q, p).
- Wigner function provides a quasi-probability distribution in phase space.
- Example: A quantum harmonic oscillator's phase space is defined using $W(q, p)$.

4.1.5 Relativity Cosmology

- General relativity: Phase space described by $(g_{\mu\nu}, \pi^{\mu\nu})$, where $g_{\mu\nu}$ is the metric and $\pi^{\mu\nu}$ its conjugate momentum.
- Cosmology: Scale factor $(a, \dot{a})$ defines the universe's evolution.
- Black hole thermodynamics: Phase space given by mass M, charge Q, and angular momentum J.

4.1.6 Quantum Field Theory String Theory

- Quantum Field Theory: Phase space defined by fields $(\phi(x), \pi(x))$.
- Gauge Theories: Uses field strengths $F_{\mu\nu}$ and conjugate momenta.
- String Theory: Uses string embedding coordinates $X^\mu(\sigma, \tau)$ and momenta $P^\mu(\sigma, \tau)$.
- Supersymmetric Theories: Extend phase space with fermionic coordinates.

4.2 Phase space of various systems



Figure 4.1: Simple Harmonic Oscillator (Elliptical Phase Space)



Figure 4.2: Damped Oscillator (Spiral Phase Space)



Figure 4.3: Nonlinear Duffing Oscillator (Strange Attractor)



Figure 4.4: Double Pendulum (Chaotic Phase Space)



Figure 4.5: Lotka-Volterra Equations (Predator-Prey Cycles)



Figure 4.6: Lorenz Attractor (Butterfly-Shaped Chaos)



Figure 4.7: KAM Tori (Kolmogorov-Arnold-Moser Theory)



Figure 4.8: Quantum Phase Space (Wigner Function)



Figure 4.9: Black Hole Phase Space (Penrose Diagram Interpretation)



Figure 4.10: Cosmic Inflation Attractors (Cosmological Phase Space)

4.3 Discretization Limit of Phase Space: Emergence of Matrix Structure

Let us consider the simplest phase space with (x, p) coordinates. To study a system, we analyze it by plotting x against p. For this purpose, we need to discretize both x and p.

Phase Space Discretization



If we want to study a planet like Earth, where the length scale is 10^7 m and the momentum scale is about 10^{29} kg·m/s, we can easily discretize phase space. By reducing the grid size further and further, we achieve higher resolution, allowing for a more precise understanding of the system.

Similarly, for a car with a length scale of 1m and a momentum scale of 10^3 kg·m/s, phase space can be discretized to analyze the system effectively.

For a dust particle with a length scale of 10^{-5} m and a momentum scale of 10^{-14} kg·m/s, further discretization enables even higher resolution for more precise study. However, the Heisenberg uncertainty principle imposes a fundamental limit, requiring that the grid area must satisfy:

$$\Delta x \Delta p \geq \frac{\hbar}{2} \approx 10^{-35} \text{ J·s}.$$

The Heisenberg uncertainty principle limits infinite resolution, i.e., the resolution of the fundamental structure of the universe is finite. This means the entire universe can be described as a discrete matrix, and $\Delta x \cdot \Delta p$ cannot be broken beyond a certain limit, which is $\Delta x \cdot \Delta p \geq \hbar/2 \approx 10^{-35}$ J·s.

If we take an analogy with an image, the size of a pixel $(\Delta x \cdot \Delta p)$ must be greater than $\hbar/2$. The universe is made of matrix elements of size $\hbar/2$.

What about an electron with a length scale of 10^{-15} m and a momentum scale of 10^{-23} kg·m/s ? If it exists within the matrix or as part of the matrix, it must satisfy $\Delta x \cdot \Delta p \geq \hbar/2$.
In order to satisfy Heisenberg's uncertainty principle (i.e., finite pixel size, which cannot be infinitely small), it must spread in either position or momentum to exist in this matrix world or as part of the matrix.

Like if you put a photon or electron in a very narrow slit (Δx is very small), the photon or electron tries to increase Δp in the x-direction, which is the diffraction phenomenon.

In general, human consciousness can perceive a length scale of approximately 1m and a momentum of 1 kg·m/s. The number of pixels or matrix elements that human consciousness perceives is given by:
$\frac{1m \times 1kg \cdot m/s}{10^{-35}} \approx 10^{35}$.
In general, a single unit of what human consciousness perceives is composed of approximately 100000000000000000000000000000000000 matrix blocks or pixels. This is akin to living in a highly detailed, high-resolution simulation.

As an analogy, increasing the pixel resolution of an image or video makes it appear more realistic. When watching a 1920 × 1080 video, we actually forget that we are viewing 2,073,600 matrix blocks or pixels, yet it appears seamless and real.

Now, comparing watching 2,073,600 matrix blocks to experiencing and perceiving more than 100000000000000000000000000000000000 matrix blocks. It becomes evident that we exist within an extraordinarily high-resolution simulation.

The discretization of phase space has a limit, indicating that phase space itself is a matrix. This concept is not restricted to (x, p) phase space but extends to all other types of phase space as well.

4.4 Absolute Vacuum or Nothingness Is a Matrix



Absolute vacuum—where no matter or field is present—is also a matrix. The phase space of absolute nothingness can be studied using energy (E) versus time (t) coordinates.

If everything were as expected and we were not part of a matrix, energy should remain zero as time evolves. However, this is not the case—energy spontaneously arises from nothing in the form of photons and other particles, momentarily, due to the Heisenberg uncertainty principle: $\Delta E \cdot \Delta t \geq \frac{\hbar}{2}$. The vacuum is a seething, dynamic environment rather than a true void.

For a time duration $\Delta t \approx \frac{\hbar}{2\Delta E}$, an energy ΔE can appear at any point in space, a phenomenon known as **Quantum Vacuum Fluctuation**. You cannot break time and energy beyond a certain limit.

We can say that everything in the universe, and the universe itself, is made of energy and time, but guess what—energy and time cannot be broken beyond a certain extent.

So, everything in this universe, including the universe itself, is a matrix, and the elements of this matrix, or pixels, are made of blocks of time and energy. The size of each time-energy block is $\Delta t \Delta E \approx \hbar/2 \approx 10^{-35}$ J·s.

4.5 List of other clues for simulation or matrix

4.5.1 Position-Momentum (x, p)

Uncertainty Relation: $\Delta x \cdot \Delta p \geq \frac{\hbar}{2}$
Associated Experiment: Electron Diffraction (Davisson-Germer)
Evidence: Wave-like diffraction confirms uncertainty.

4.5.2 Energy-Time (t, E)

Uncertainty Relation: $\Delta E \cdot \Delta t \geq \frac{\hbar}{2}$
Associated Experiment: Decay of Unstable Particles
Evidence: Short-lived Particles have broader energy distributions.

4.5.3 Angular Position-Angular Momentum (θ, L_z)

Uncertainty Relation: $\Delta \theta \cdot \Delta L_z \geq \frac{\hbar}{2}$
Associated Experiment: Optical Vortex Beams
Evidence: Angular localization increases momentum uncertainty.

4.5.4 Number-Phase (N, ϕ)

Uncertainty Relation: $\Delta N \cdot \Delta \phi \geq \frac{1}{2}$
Associated Experiment: Squeezed Light Experiments
Evidence: Phase squeezing reduces $\Delta \phi$ while increasing ΔN.

Chapter 5

Video Game vs. Our Universe

5.1 4K Video Game: A Simulation Created by Humans

5.1.1 Resolution or Size of the Matrix

Resolution:

$$3840 \times 2160 = 8,294,400 \text{ pixels} \approx 10^6 \text{ pixels}$$

5.1.2 How the Image Matrix Changes Over Time

A computer program, written in binary (1 for "on" and 0 for "off"), simulates physical reality using billions of transistors on a graphics card, which is optimized for matrix operations.

5.1.3 Powered By

Electrical Energy

5.2 Our Universe

5.2.1 Resolution or Size of the Matrix

Let's calculate the resolution of the universe in (x, p) Coordinates

1. **Range of** x: The spatial coordinate x can range from 0 to the diameter of the universe. Since the exact diameter is unknown, we use the observable universe's diameter, approximately 10^{26} m.
    - *Note:* We are only making a rough estimate of how many times the universe has a higher resolution than a 4K video game.
2. **Range of** p: The momentum p can range from 0 to a large value. For estimation, we take the momentum of the Milky Way galaxy, approximately 10^{48} kg·m/s.

- We could choose an even larger momentum, but this is sufficient for a rough comparison.

3. **Calculating Resolution**: The resolution of phase space can be estimated as:

$$\text{Resolution} \gtrsim \frac{x_{\max} \cdot p_{\max}}{\Delta x \cdot \Delta p} \tag{5.1}$$

Given:

$$\begin{aligned} x_{\max} &\approx 10^{26} \text{ m} \\ p_{\max} &\approx 10^{48} \text{ kg·m/s} \\ \Delta x \cdot \Delta p &\approx 10^{-35} \text{ J·s} \end{aligned}$$

$$\begin{aligned} \text{Resolution} &\gtrsim \frac{10^{26} \times 10^{48}}{10^{-35}} \\ &\gtrsim 10^{109} \end{aligned}$$

= 10000000000

1. **Second Law of Thermodynamics**:

$$S \geq 0 \tag{5.3}$$

Entropy of an isolated system never decreases, ensuring the arrow of time.

2. **Principle of Least Action**:

$$\delta S = 0, \quad S = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt$$

A system chooses a stationary action path to transition from one state to another.

3. **Maxwell's Equations**:

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}, \quad \nabla \cdot \mathbf{B} = 0 \tag{5.4}$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \quad \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \tag{5.5}$$

Describe how electric and magnetic fields interact and evolve.

4. **Schrödinger's Equation**:

$$i\hbar \frac{\partial}{\partial t} \Psi = \hat{H} \Psi \tag{5.6}$$

Governs the quantum state evolution of a system.

5. **Boltzmann Entropy Formula**:

$$S = k_B \ln \Omega \tag{5.7}$$

Connects entropy to the number of microscopic states.

6. **Relativistic Energy-Momentum Relation**:

$$E^2 = p^2 c^2 + m^2 c^4 \tag{5.8}$$

Relates energy, momentum, and mass in special relativity.

7. **Einstein Field Equations**:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \tag{5.9}$$

Describe how matter and energy shape spacetime curvature.

8. **Dirac Equation**:

$$(i\gamma^\mu \partial_\mu - m)\psi = 0 \tag{5.10}$$

Describes relativistic quantum particles with spin-$\frac{1}{2}$.

9. **Feynman Path Integral**:

$$\langle x_f | e^{-iHt/\hbar} | x_i \rangle = \int \mathcal{D}[x] e^{iS[x]/\hbar} \tag{5.11}$$

Reformulates quantum mechanics by summing over all possible paths a particle can take.

10. **Yang-Mills Equations (Gauge Theories)**:

$$D_\mu F^{\mu\nu} = J^\nu \tag{5.12}$$

Generalize Maxwell's equations to non-Abelian gauge fields.

11. **Hawking Radiation Equation**:

$$T_H = \frac{\hbar c^3}{8\pi G M k_B} \tag{5.13}$$

Predicts black holes emit thermal radiation due to quantum effects.

12. **Friedmann Equations**:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho - \frac{k}{a^2} \tag{5.14}$$

Describe the expansion dynamics of the universe.

5.2.3 Powered By

Energy of different forms: gravitational, electromagnetic, nuclear, and dark energy. Additional forms of energy may exist but have yet to be discovered.